

PostOp Instructions

Bone Grafting

Your bone graft is made up of many particles. You may find some small granules in your mouth for the first several days. Do not be alarmed by these. The socket is over packed to allow for the loss of a small amount of particles. It is normal to have some of them come out of the graft site and into your mouth. There are some things that can be done to minimize the amount of particles that become dislodged.

What to expect following surgery:

BLEEDING: Small amounts of blood in the saliva can make your saliva appear quite red/pink the remainder of the day after the procedure.

PAIN: Before the anesthetic wears off it is recommended that you fill your prescriptions given to you at the time of the surgery. **If more severe discomfort occurs please contact the office 626-304-3004.**

SWELLING: Some swelling of the lip or cheek may occur.

What to do following surgery:

After leaving the office, rest and avoid strenuous activities for the remainder of the day. This will help reduce bleeding and aid in the healing process.

Take your meds as described above. Be sure to take before the anesthesia wears off. Nausea can occur and most often is due to taking your meds on an empty stomach. Take your meds with a large glass of water or with some soft food.

Eat soft foods for the first 2-4 days. Maintain a good, balanced diet. Drink plenty of water. Avoid chewing hard foods on the graft sites. Chewing forces during the healing phase can decrease the body's ability to heal around the graft.

Do not use a straw. Avoid alcohol for 48 hours. For the first day, it is advisable to let the blood clot stabilize by not rinsing your mouth.

Following the first day, gentle rinsing would be advised; avoid vigorous rinsing. Do not rinse vigorously because you can disturb some of the bone graft granules. Do not use any mouthwash. Use only water or warm salt water if desired. Smoking should be avoided for a minimum of 1 week as it slows down the healing process.

Do not apply pressure with your tongue or fingers to the grafted area because the material is movable during the initial healing. Do not lift or pull the lip to look at the sutures as this can actually damage the site or tear the sutures and cause damage to the wound site. Avoid poking the site with your tongue.

Do not avoid cleaning the area. You can take a q-tip and dip it in hydrogen peroxide or just water and gently clean the site. Do not avoid brushing your teeth but stay away from the site with the toothbrush. Let comfort be your guide.

Please call the office if you have: Uncontrollable pain - Excessive or severe bleeding - Marked Fever - Excessive warm swelling. **Or if you have any other questions or concerns please contact our office at (626)304-3004**